


Point-of-Care Test Developed for Tumor Marker in Human Saliva Based on Lanthanide Nanoprobes



2021-01-25

Salivary assay, emerging as a non-invasive alternative to blood assay in clinic analysis, holds great promise for early-stage cancer diagnostics with advantages of low cost, easy collection and facile processing. Therefore, point-of-care (POC) detection of tumor markers in the saliva is urgently demanded.

However, the salivary assay has been severely hindered by the limitation of inadequate sensitivity in current commercial assay kits because of the much lower level of tumor markers in saliva than in human serum.

In a study published in [Advanced Science](#), a research group led by Prof. CHEN Xueyuan from [Fujian Institute of Research on the Structure of Matter \(FJIRSM\)](#) of the [Chinese Academy of Sciences](#) developed a unique lab-in-syringe strategy for rapid and ultrasensitive detection of tumor markers in saliva based on lanthanide nanoprobes.

The researchers employed Eu_2O_3 nanocrystals as bioprobes, which can be easily dissolved in acidic enhancer solution and transformed into a large number of highly luminescent Eu^{3+} micelles. Meanwhile, they utilized a disposable syringe filter equipped with nitrocellulose membrane as bioassay platform, which facilitated the accomplishment of detection process within 10 min.

By ingenious integration of dissolution enhanced luminescent bioassay strategy and the miniaturized detection device, the researchers demonstrated the feasibility and reliability for the direct quantitation of tumor marker like carcinoembryonic antigen (CEA) in patient saliva samples with a detection limit down to 1.47 pg/mL (7.35 fM).

More importantly, by virtue of such an excellent luminescence-amplification strategy, the researchers can visually detect the photoluminescence intensity change above 0.1 ng/mL (0.5 pM) of CEA by naked eyes to qualitatively evaluate its level in saliva. The whole detection process is easy to operate, which is highly beneficial for cancer diagnostics by ordinary people at home.

These findings revealed the great potential of the proposed general strategy in practical home self-monitoring of trace amount of tumor markers in saliva, which may accelerate the exploitation of lanthanide nanoprobe for POC bioassay of diverse disease markers in complex biological fluids.

Read the [original article](#) on Chinese Academy of Sciences.