
Study Reveals Mechanism of Ag Nanoclusters Dispersion Induced by Oxygen

2021-03-11

Oxidative dispersion has been widely used in the regeneration of sintered metal catalysts as well as the fabrication of single-atom catalysts.

The oxidative dispersion process includes the formation of mobile metal oxide species from large metal particles and the capture of these species on a support surface. Nevertheless, the mechanism of oxidation-induced dispersion has yet to be confirmed via in situ electron microscopic and/or spectroscopic characterizations.

Recently, a research team led by Prof. FU Qiang and Prof. BAO Xinhe from the Dalian Institute of Chemical Physics (DICP) of the Chinese Academy of Sciences ([CAS](#)), in collaboration with Prof. YANG Bing from DICP and Prof. GAO Yi from the Shanghai Institute of Applied Physics of CAS, reported the oxygen adsorption-induced dispersion of metallic Ag nanoclusters in typical oxidative condition.

The results were published in [Nature Communications](#) on March 3.



Dynamic evolution of Ag nanostructures during oxidative dispersion.

By utilizing in situ imaging methods such as environmental scanning electron microscopy (ESEM), and newly developed near-ambient pressure photoemission electron microscopy (NAP-PEEM), the researchers found that micron-scale Ag nanowires could be dispersed into subnanometer clusters under oxygen condition.

Ex situ experiments indicated that Ag nanowires were converted into AgOx nanoclusters.

Conversely, in this study, in situ near-ambient pressure photoelectron spectroscopy (NAP-XPS) directly demonstrated the presence of a transitional state of metallic Ag nanoclusters during dispersion at high temperatures, while the formation of the oxide occurred during the cooling process. The dynamic dispersion of Ag nanowires during CO oxidation was also demonstrated.

Experimental and theoretical calculations showed that chemisorption of oxygen from the O₂ condition was the essential driving force for the dispersion of metallic Ag nanoclusters.

This work provides a new understanding of the role of the O₂ condition in oxidative dispersion, which is particularly important for the prediction and control of the dynamic dispersion/redispersion of supported metal catalysts under similar reaction conditions.

Read the [original article](#) on Chinese Academy of Sciences (CAS).