

Exploring the Nanoworld in 3D

2021-04-02

On the nanometric scale (one billionth of a metre), materials can feature new properties. A French team together with an Austrian one recently took an important step forward by mapping one of these properties for the first time in 3D.

Imagine a cube on which light is projected by a flashlight. The cube reflects the light in a particular way, so simply spinning the cube or moving the flashlight makes it possible to examine each aspect and deduce information regarding its structure. Now, imagine that this cube is just a few atoms high, that the light is detectable only in infrared, and that the flashlight is a beam from a microscope. How to go about examining each of the cube's sides? That is the question recently answered by scientists from the CNRS, [l'Université Paris-Saclay](#), the [University of Graz](#) and [Graz University of Technology \(Austria\)](#) by generating the first 3D image of the structure of the infrared light near the nanocube. Their results will be published on 26 March 2021 in [Science](#).

Electron microscopy uses an electron beam to illuminate a sample and create an enlarged image. It also provides more complete measurements of physical properties, with unrivalled spatial resolution that can even visualize individual atoms. Chromatem, the Equipex Tempos team's dedicated instrument for spectroscopy, is one of these new generation microscopes. It probes the optical, mechanical, and magnetic properties of matter with very high resolution, one that is matched by only three other microscopes in the world.

Scientists from the CNRS and l'Université Paris-Saclay working at the Solid States Physics Laboratory (CNRS/Université Paris-Saclay), along with their colleagues at the University of Graz and Graz University of Technology ([Austria](#)), used Chromatem to study a magnesium oxide nanocrystal. The vibration of its atoms creates an electromagnetic field that can only be detected in the mid-infrared range[2]. When the electrons emitted by the microscope indirectly encounter this electromagnetic field, they lose energy. By measuring this energy

loss, it becomes possible to deduce the outlines of the electromagnetic field surrounding the crystal. The problem is that this type of microscopy can only provide images in 2D, raising the question of how to visualize all of the cube's corners, edges, and sides. In order to do so, the scientists developed image reconstruction techniques that have, for the first time, generated 3D images of the field surrounding the crystal. This will eventually enable targeting a specific point on the crystal, and conducting localized heat transfers, for instance.

Many other nano-objects absorb infrared light, such as during heat transfers, and it will now be possible to provide 3D images of these transfers. This is one avenue of exploration for optimizing heat dissipation in the increasingly small components used in nanoelectronics.

Read the [original article](#) on EurekAlert.