

SARS-CoV-2 Cytokine Storm Inhibited by Curcumin Nanoparticles



2022-02-12

In a recent study published in the journal ACS Applied Bio Materials, researchers demonstrated the inhibitory potential of curcumin-encapsulated polysaccharide nanoparticles (CUR–PS-NPs) against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) spike protein (CoV2-SP)-stimulated cytokine storm in lung and liver epithelial cells.

The multi-organ failure, lung injury, and poor prognosis of severe disease in coronavirus disease 2019 (COVID-19) patients are directly associated with the interleukin (IL)-induced profound cytokine storm developed in response to SARS-CoV-2 infection in these patients. The interaction between SARS-CoV-2 and human angiotensin-converting enzyme 2 (hACE2) receptors triggers the cytokine storm followed by life-threatening systemic inflammatory responses. The fatal inflammatory syndrome is characterized by higher circulating cytokine levels and immune cell hyperactivation, leading to secondary organ dysfunction, especially hepatic, pulmonary, or renal.

A dietary polyphenolic antioxidant compound named CUR obtained from turmeric plant *Curcuma longa* L. is famous for its robust anti-inflammatory activity. However, in vivo efficacy of CUR is constrained because of its inadequate bioavailability.

About the study

In the present [study](#), the authors compared the anti-inflammatory effect of CUR–PS-NPs directed towards CoV2-SP-stimulated cytokine storm with bulk-CUR (B-CUR) and PS-NPs in lung and liver epithelial cells. Further, the researchers interpreted the mechanism associated with the inhibition of nuclear factor κ B (NF- κ B) signaling and mitogen-activated protein kinase (MAPK) deactivation by CUR that inhibited the CoV2-SP-mediated cytokine storms. The team also evaluated the function of CUR in lowering CoV2-SP-induced phosphorylation of p42/44 MAPK, p65/NF- κ B, p38 MAPK, and nuclear p65/NF- κ B expression, and release of growth factors, chemokines, and cytokines associated with the lung and liver epithelial cell injury.

CUR (97% purity), purified biotinylated recombinant hACE2, and SARS-CoV-2 S recombinant glycoproteins were used in this study. Further, the liver Huh7.5 and lung A549 epithelial cells were procured from the American Type Culture Collection ([ATCC](#))-recognized cell repository at [National Centre for Cell Sciences](#), Pune, [India](#).

The PS-rich fraction was isolated and characterized from the dried rhizomes of *C. longa*. An emulsion solvent evaporation technique with minor changes was used in the CUR–PS-NPs' preparation. The epithelial cells were seeded and incubated with CoV2-SP containing stock solution.

The total ribonucleic acid (RNA) was isolated from the CoV2-SP-exposed epithelial cell pallets using an RNeasy mini kit. Subsequently, reverse transcription-quantitative polymerase chain reaction (RT-qPCR) experiments were conducted in the genes of CoV2-SP-exposed and unexposed epithelial cells. The interaction between CoV2-SP and hACE2 was assessed by enzyme-linked immunosorbent assay (ELISA). The cell viability was evaluated by the fluorescence-activated cell sorting (FACS) analysis, fluorescence microscopy, and Alamar blue assay using propidium iodide (PI) and Annexin V-FITC.

Study findings

The results indicated that the size of the CUR–PS-NPs was within the nanomaterial range, and there was a continuous CUR release from the CUR–PS-NPs.

CUR–PS-NPs showed a high potential to inhibit the release of chemokines such as IP10, MCP-1, and CXCL8, growth factors such as vascular endothelial growth factor (VEGF) and C-C chemokine ligand 3 (CCL3), and cytokines such as IL-12, IL-6, and IL-1 associated with the impairment of CoV2-SP-triggered lung A549 and liver Huh7.5 epithelial cells.

The inhibition of chemokines, cytokines, and growth factors by CUR–PS-NPs in A549 and Huh7.5 cells was higher than PS-NPs and B-CUR-treated cells. However, the inhibition of VEGF-A, CCL10, and CCL5 in PS-NPs and B-CUR-treated A549 cells were higher than the CUR–PS-NPs-treated cells. No significant inhibitory effects on the release of IL-8 and IL-6 in CoV2-SPs-triggered Huh7.5 and A549 cells were demonstrated by the PS-NPs treatment alone.

Therapeutical strategies employing CUR-PS-NPs successfully diminished the interaction between CoV2-SP and host ACE2. CUR-PS-NPs' potential to inhibit the interaction of CoV2-SP and human ACE2 was higher than that of PS-NPs and B-CUR.

At the 10 μ M dose, both B-CUR and nano-CUR caused significant reductions in cell viability whereas, at the 5 μ M dose, neither B-CUR nor nano-CUR caused any lethality to the cells. Additionally, nano-CUR had superior bioavailability than B-CUR.

CUR-PS-NPs act by lowering NF- κ B/MAPK signaling, resulting in the reduction of CoV2-SP-induced phosphorylation of p65/NF- κ B, p38 MAPK, p65/NF- κ B, and p42/44 MAPK expression. The reduction of CoV2-SP-induced phosphorylation of protein kinases caused by CUR-PS-NPs was higher than B-CUR-induced reductions.

Conclusions

The study findings indicated that organic NPs of CUR obtained from a PS-affluent fraction of turmeric rhizome, CUR-PS-NPs, have a high inhibitory potential towards CoV2-SP-induced cytokine storm in human lung A549 and liver Huh7.5 epithelial cells. Furthermore, the treatment of CUR-PS-NPs in CoV2-SP-triggered epithelial cells prevented the epithelial cell damage by inhibiting the release of chemokines, growth factors, and cytokines through the deactivation of the NF- κ B/MAPK signaling pathway.

Overall, the study emphasized that the CUR-PS-NPs might have the potential to become a safe and effective drug for COVID-19 treatment. However, additional *in vivo* investigations are needed to validate the CoV2-SP-induced cytokine storm inhibitory potential of CUR-PS-NPs in lung and liver epithelial cells.

Read the [original article](#) on News Medical.